

APPROXIMATION OF A FUNCTION BELONGING TO  $LIP(\alpha, \theta, \omega)$  BY  $(C, 1)(E, q)$  MEAN

SMITA SONKER<sup>1</sup>, ALKA MUNJAL<sup>2</sup> AND LAKSHMI NARAYAN MISHRA<sup>\*,3</sup>

<sup>1</sup> Department of Mathematics, N.I.T. Kurukshetra, Haryana

<sup>2</sup> Department of Mathematics, Akal University, Bathinda, Punjab

<sup>3</sup> Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology (VIT) University, Vellore 632 014, Tamil Nadu, India

*smita.sonker@gmail.com, alka\_math@auts.ac.in,*

*Corresponding Author: \*lakshminarayanmishra04@gmail.com*

**Abstract:** A result has been established for a conjugate trigonometric Fourier series in which degree of approximation determine in class  $Lip(\alpha, \theta, w)$  by a product  $(C, 1)(E, q)$  mean (sequence to sequence transformation). This result is more applicable by using the product mean as it is used as double filter.

**Key Words:** Signal approximation;  $(C, 1)(E, q)$  mean; conjugate Fourier series; Hölder's inequality; Minkowski's inequality.

**AMS (MOS) Subject Classification.** 40G05, 42B05, 42B08, 41A25.

1. INTRODUCTION

Let  $\{s_n\}$  represents the partial sum sequence of series  $\sum a_n$ . The  $(C, 1)$  transformed mean of  $\{s_n\}$  is

$$(1.1) \quad C_n^1 = \frac{1}{n+1} \sum_{k=0}^n s_k, \quad n = 0, 1, 2, \dots$$

It is said to be Cesàro summable of order one  $(C, 1)$  to  $s$  if  $\lim_{n \rightarrow \infty} C_n^1 = s$ . Euler transformed means of  $q > 0$  of  $\{s_n\}$  is

$$(1.2) \quad E_n^q = \frac{1}{(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} s_k, \quad q > 0, n = 0, 1, 2, \dots$$

Let the product  $(C, 1)(E, q)$  transformed mean denoted by  $C_n^1 E_n^q$  and given by

$$(1.3) \quad C_n^1 E_n^q = \frac{1}{n+1} \sum_{k=0}^n E_k^q = (1+n)^{-1} \sum_{k=0}^n (1+q)^{-k} \sum_{v=0}^k \binom{k}{v} q^{k-v} s_v.$$

Let  $f$  is  $2\pi$ -periodic function and belongs to  $L_\theta[0, 2\pi] = L_\theta$  for  $\theta \geq 1$  (bounded, integrable). Then,

$$(1.4) \quad s_n(f; x) = \frac{a_0}{2} + \sum_{l=1}^n (a_l \cos lx + b_l \sin lx)$$

and modulus of continuity of function  $f$  is

$$(1.5) \quad \Omega_\theta(\delta; f) = \sup_{0 < |h| \leq \delta} \left\{ \frac{1}{2\pi} \int_0^{2\pi} |f(x+h) - f(x)|^\theta dx \right\}^{1/\theta}.$$

If, for  $\alpha > 0$ , then condition for  $f \in Lip(\alpha, \theta)$ ,  $\theta \geq 1$  is

$$(1.6) \quad \Omega_\theta(\delta; f) = O(\delta^\alpha).$$

The  $L_\theta$ -norm of the function is

$$(1.7) \quad \|f\|_\theta := \left\{ \frac{1}{2\pi} \int_0^{2\pi} |f(x)|^\theta dx \right\}^{1/\theta}.$$

For the function  $f$ , conjugate Fourier series is

$$(1.8) \quad \sum_{l=1}^{\infty} (-a_l \sin lx + b_l \cos lx)$$

and

$$(1.9) \quad \tilde{s}_n(f; x) = \sum_{l=1}^n (-a_l \sin lx + b_l \cos lx)$$

represents the partial sum. If  $\psi(t) = f(x+t) - f(x-t)$  and conjugate  $\tilde{f}$  is

$$(1.10) \quad 2\pi\tilde{f}(x) = -\lim_{\varepsilon \rightarrow 0} \int_\varepsilon^\pi \cot(t/2)\psi(t)dt.$$

If  $f \in Lip \alpha$ , then condition is

$$(1.11) \quad |f(x+t) - f(x)| = O(t^\alpha).$$

If  $0 < \alpha \leq 1$  and  $\theta \geq 1$ , then  $\left( \int_0^{2\pi} |f(x+t) - f(x)|^\theta dx \right)^{1/\theta} = O(t^\alpha)$  is condition for  $f \in Lip(\alpha, \theta)$ .

For  $r \geq 1$ , the  $L_\theta$ -norm for  $f \in L_\theta[-\pi, \pi]$  is

$$(1.12) \quad \|f\|_\theta = \left( \int_0^{2\pi} |f(x)|^\theta dx \right)^{1/\theta}.$$

For  $\theta \geq 1$ , the condition for the function  $f \in L_w^\theta[0, 2\pi] = L_w^\theta$  is

$$(1.13) \quad \|f\|_{\theta, w} = \left( \int_0^{2\pi} w(x)|f(x)|^\theta dx \right)^{1/\theta} < \infty.$$

If

$$(1.14) \quad \sup_I \left( \frac{1}{|I|} \int_I [w(x)]^{-1/(\theta-1)} dx \right)^{\theta-1} \left( \frac{1}{|I|} \int_I w(x) dx \right) < \infty, \quad \theta > 1.$$

Then, function belongs to  $A_\theta$  i.e. Muckenhoupt class, where the supremum is in  $|I| \leq 2\pi$ .

Let  $w \in A_\theta$  and  $f \in L_w^\theta$ . The condition

$$(1.15) \quad \Omega_\theta(f; \delta) = \sup_{0 < |h| \leq \delta} \|\Delta_h(f)\|_{\theta, w}$$

represents the modulus of continuity, where

$$(1.16) \quad \Delta_h(f)(x) = \frac{1}{h} \int_0^h |f(x+t) - f(x)| dt.$$

The  $Lip(\alpha, \theta, w)$  for  $0 < \alpha \leq 1$  and  $\delta > 0$  is

$$(1.17) \quad Lip(\alpha, \theta, w) = \{f \in L_w^\theta : \Omega(f, \delta)_{\theta, w} = O(\delta^\alpha)\}.$$

The problem in which approximation of a signal can done solved by selecting a function closely matches (approximates) a target function among a well-defined class. Quade [17] solved a problem related with approximation by trigonometric polynomials. Many research articles have been devoted to the study of summability of infinite series due to its wide range of applications. After this, several mathematicians determined various results on approximation by summability techniques of a signal which belongs to various classes like Chandra [1-3], Khan [4, 5], Leindler [6], Mittal et al. [8] and Mishra et al. [9-16]. Recently, Sonker and Munjal [18-25] gave a number of theorems exploring summability of the Orthogonal, Fourier series and infinite series. In the present result, summability method has been used for sharper estimations.

## 2. KNOWN RESULTS

A result on  $(C, 1)(E, 1)$  transformed means has been established by Lal and Singh [7].

**Theorem 1** [7]: A function belonging to  $Lip(\alpha, r)$  and  $2\pi$ -periodic ( $f : R \rightarrow R$ ), then by the  $(C, 1)(E, 1)$  transformed mean of conjugate Fourier series, the degree of approximation (doa) of  $\tilde{f}(x)$  of  $f$  satisfies,

$$(2.1) \quad M_n(f) = Min \left\| (CE)_n^1 - \tilde{f} \right\|_r = O(n^{1/r-\alpha}), \quad n = 0, 1, 2, \dots$$

and  $(C, 1)(E, 1)$  transformed mean is

$$(2.2) \quad (CE)_n^1 = (n+1)^{-1} \sum_{k=0}^n \left( 2^{-k} \sum_{i=0}^k \binom{k}{i} S_i \right).$$

## 3. MAIN RESULT

In this paper, we establish a result for a function belonging to  $Lip(\alpha, \theta, w)$  by  $(C, 1)(E, q)$  mean with the help of conjugate trigonometric Fourier series.

**Theorem 2:** Let function  $f \in Lip(\alpha, \theta, w)$ -class with  $\theta \geq 1$ ,  $\alpha\theta \geq 1$  which is Lebesgue integrable  $2\pi$ -periodic, then degree of approximation of  $\tilde{f}(x)$  with  $(C, 1)(E, q)$  transformed means of conjugate Fourier series of  $f$  satisfies,

$$(3.1) \quad M_n(f) = Min \left\| (C_n^1 E_n^q) - \tilde{f} \right\|_{\theta, w} = O(n^{\frac{1}{\theta}-\alpha}), \quad \text{where } n = 0, 1, 2, \dots,$$

provided

$$(3.2) \quad \left( \int_0^{\pi/(n+1)} (|\psi(t)| t^{-\alpha})^\theta \right)^{1/\theta} = O\left(\frac{1}{n+1}\right)$$

$$(3.3) \quad \left( \int_0^{\pi/(n+1)} (|\psi(t)| t^{-\delta-\alpha})^\theta \right)^{1/\theta} = O(n+1)^\delta,$$

where  $\delta$  is an arbitrary number with  $1/\theta + 1/\varphi = 1$  and  $(\alpha + \delta)\varphi + 1 < 0$  for  $\theta > 1$ .

#### 4. LEMMAS

In order to prove our theorem, we need the following lemmas.

**Lemma 1 [7]:**  $|K_n(t)| = O((n+1)t) + O(1/t)$  for  $\pi/(n+1) \geq t \geq 0$ .

**Lemma 2 [7]:**  $|K_n(t)| = O(1) + O(1/t)$  for  $\pi/(n+1) \leq t \leq \pi$ .

#### 5. PROOF OF MAIN THEOREM

The  $\tilde{s}_n(f; x)$  is represented by

$$(5.1) \quad \tilde{s}_n(f; x) = \frac{1}{\pi} \int_0^\pi \frac{\cos(n+1/2)t - \cos(t/2)}{2 \sin(t/2)} \psi(t) dt.$$

Therefore,

$$(5.2) \quad \tilde{f}(x) - \tilde{s}_n(f; x) = -\frac{1}{2\pi} \int_0^\pi \frac{\cos(n+1/2)t}{\sin(t/2)} \psi(t) dt.$$

The  $(C, 1)(E, q)$  transform

$$(5.3) \quad \begin{aligned} \tilde{f} - C_n^1 E_n^q &= -\frac{1}{2\pi(n+1)} \left[ \sum_{k=0}^n (1+q)^{-k} \int_0^\pi \frac{\psi(t)}{\sin(t/2)} \sum_{v=0}^k \binom{k}{v} q^{k-v} \cos[(v+1/2)t] dt \right] \\ &= \left[ \int_0^{\frac{\pi}{n+1}} + \int_{\frac{\pi}{n+1}}^\pi \right] \psi(t) K_n(t) dt = I_1 + I_2 \quad (\text{say}). \end{aligned}$$

Using condition (3.2) and Lemma 1, we have

$$(5.4) \quad \begin{aligned} |I_1| &= \int_0^{\pi/(n+1)} |K_n(t)| |\psi(t)| dt \leq \left[ \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^{\pi/(n+1)} (|K_n(t)| t^\alpha)^\varphi dt \right]^{1/\varphi} \left[ \int_0^{\pi/(n+1)} \left( \frac{|\psi(t)|}{t^\alpha} \right)^\theta dt \right]^{1/\theta} \\ &= O((n+1)^{-1}) \left[ \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^{\pi/(n+1)} ((n+1)t^{\alpha+1} + t^{\alpha-1})^\varphi dt \right]^{1/\varphi} \\ &= O((n+1)^{-1}) \left[ \left( \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^{\pi/(n+1)} (n+1)t^{(\alpha+1)\varphi} dt \right)^{1/\varphi} + \left( \lim_{\varepsilon \rightarrow 0} \int_\varepsilon^{\pi/(n+1)} t^{(\alpha-1)\varphi} dt \right)^{1/\varphi} \right] \\ &= O((n+1)^{-1}) \left[ (n+1)(n+1)^{-\alpha-1-1/\varphi} + (n+1)^{-\alpha+1-1/\varphi} \right] \\ &= O((n+1)^{-1}) \left[ (n+1)(n+1)^{-\alpha-1+1/\theta} + (n+1)^{-\alpha+1/\theta} \right] \\ &= O \left[ (n+1)(n+1)^{-\alpha-2+1/\theta} + (n+1)^{-\alpha-1+1/\theta} \right] \\ &= O \left( (n+1)^{-\alpha-1+1/\theta} \right). \end{aligned}$$

Now, we consider and using condition (3.3) and

$$\begin{aligned} &= O((n+1)^\delta) \left[ \left( \int_{\pi/(n+1)}^\pi t^{(\alpha+\delta)\varphi} dt \right)^{1/\varphi} + \left( \int_{\pi/(n+1)}^\pi t^{(\alpha+\delta-1)\varphi} dt \right)^{1/\varphi} \right] \\ &= O((n+1)^\delta) \left[ (n+1)^{-\alpha-\delta-1/\varphi} + (n+1)^{-\alpha-\delta+1-1/\varphi} \right] \\ &= O \left[ (n+1)^{-\alpha-1/\varphi} + (n+1)^{-\alpha+1-1/\varphi} \right] \end{aligned}$$

$$\begin{aligned}
&= O\left[(n+1)^{-\alpha-1+1/\theta} + (n+1)^{-\alpha+1/\theta}\right] \\
(5.5) \quad &= O\left((n+1)^{-\alpha+1/\theta}\right).
\end{aligned}$$

Combining (5.1)-(5.5), we have

$$(5.6) \quad \left|C_n^1 E_n^q - \tilde{f}\right| = O\left((n+1)^{1/\theta-\alpha}\right)$$

Hence,

$$(5.7) \quad \left\|C_n^1 E_n^q - \tilde{f}\right\|_{\theta, w} = O\left(\int_0^{2\pi} \left|C_n^1 E_n^q - \tilde{f}(x)\right|^\theta w(x) dx\right)^{1/\theta} = O\left((n+1)^{1/\theta-\alpha}\right)$$

This completes the proof.

**Remark:** The result will be for  $(C, 1)(E, 1)$  means, if the result can be found by using  $q=1$  in  $(C, 1)(E, q)$  transformation.

## 6. Corollaries

**Corollary 1:** If the function belong to  $\text{Lip}\alpha$ ,  $0 < \alpha \leq 1$  class which is Lebesgue integrable function  $f : R \rightarrow R$  and  $2\pi$ -periodic, then the degree of approximation of  $\tilde{f}(x)$  of conjugate Fourier series of  $f$  by the  $(C, 1)(E, q)$  transformed means satisfies,

$$\left\|(C_n^1 E_n^q) - \tilde{f}\right\|_\infty = O\left(\frac{1}{n^\alpha}\right), \quad n = 0, 1, 2, \dots$$

**Proof.** For proof we take  $w(x)=1$  and  $\theta \rightarrow \infty$  in the main theorem which reduces to the corollary 1.

## ACKNOWLEDGMENT

This work has been financially supported by Science and Engineering Research Board (SERB) through Project No.: EEQ/2018/000393. The authors offer their true thanks to the Science and Engineering Research Board for giving financial support.

## REFERENCES

- [1] P. Chandra, Trigonometric approximation of functions in  $L_p$ -norm, *J. Math. Anal. and Math. Appl.*, vol. 275, pp. 13-26, 2002.
- [2] P. Chandra, On the degree of approximation of a class of functions by means of Fourier series, *Acta Mathematica Hungarica*, vol. 52, no. 3-4, pp. 199-205, 1988.
- [3] P. Chandra, A note on the degree of approximation of continuous functions, *Acta Mathematica Hungarica*, vol. 62, no. 1-2, pp. 21-23, 1993.
- [4] H. H. Khan, On the degree of approximation of a functions belonging to the class  $\text{Lip}(\alpha, p)$ , *Indian Journal of Pure and Applied Mathematics*, vol. 5, pp. 132-136, 1974.
- [5] H. H. Khan, Approximation of classes of functions [Ph.D. thesis], *AMU, Aligarh*, India, 1974.
- [6] L. Leindler, Trigonometric approximation of functions in  $L_p$  norm, *Mathematical Analysis and Application*, vol. 302, no. 1, pp. 129-136, 2005.
- [7] S. Lal, P. N. Singh, Degree of approximation of conjugate of  $\text{Lip}(\alpha, p)$  function by  $(C, 1)(E, 1)$  means of conjugate series of a Fourier series, *Tamkang J. Math.*, vol. 33, no. 3, pp. 269-274, 2002.
- [8] M. L. Mittal, B. E. Rhoades, S. Sonker, U. Singh, Approximation of signals of class  $\text{Lip}(\alpha, p)$  by linear operator, *Applied Mathematics and Computation*, vol. 217, no. 9, pp. 4483-4489, 2011.

- [9] L. N. Mishra, P. K. Das, P. Samanta, M. Misra, U. K. Misra, On indexed absolute matrix summability of an infinite series, *Applications and Applied Mathematics*, vol. 13, 274-285, 2018.
- [10] L. N. Mishra, D. Acharya, S. Sahu, P. C. Nayak, U. K. Misra, Indexed absolute summability factor of improper integrals, *Applications and Applied Mathematics*, vol. 15, no. 1, pp. 666-672, 2020.
- [11] L. N. Mishra, M. Patro, S. K. Paikray, B. B. Jena, A certain class of statistical deferred weighted  $\mathcal{A}$ -summability based on  $(p, q)$ -integers and associated approximation theorems, *Applications and Applied Mathematics*, vol. 14, no. 2, pp. 716-740, 2019.
- [12] V. N. Mishra, Some Problems on Approximations of Functions in Banach Spaces, Ph.D. Thesis, *Indian Institute of Technology, Roorkee* 247 667, Uttarakhand, India, 2007.
- [13] V. N. Mishra, L. N. Mishra, Trigonometric Approximation of Signals (Functions) in  $L_p$ -norm, *International Journal of Contemporary Mathematical Sciences*, vol. 7, no. 19, pp. 909-918, 2012.
- [14] L. N. Mishra, V. N. Mishra, K. Khatri, Deepmala, On The Trigonometric approximation of signals belonging to generalized weighted Lipschitz  $W(L^r, \xi(t)) (r \geq 1)$ - class by matrix  $(C^1.N_p)$  Operator of conjugate series of its Fourier series, *Applied Mathematics and Computation*, vol. 237, pp. 252-263, 2014.
- [15] V. N. Mishra, K. Khatri, L. N. Mishra, Deepmala, Trigonometric approximation of periodic Signals belonging to generalized weighted Lipschitz  $W'(L_r, \xi(t)) (r \geq 1)$ - class by Nörlund-Euler  $(N, p_n)(E, q)$  operator of conjugate series of its Fourier series, *Journal of Classical Analysis*, vol. 5, no. 2, pp. 91-105, 2014.
- [16] Deepmala, L. N. Mishra, V. N. Mishra, Trigonometric Approximation of Signals (Functions) belonging to the  $W(L_r, \xi(t)) (r \geq 1)$ - class by  $(E, q)$  ( $q > 0$ )-means of the conjugate series of its Fourier series, *GJMS Special Issue for Recent Advances in Mathematical Sciences and Applications-13, Global Journal of Mathematical Sciences*, vol. 2, no. 2, pp. 61-69, 2014.
- [17] E. S. Quade, Trigonometric approximation in mean, *Duke Math. J.* vol. 3, pp. 529-542, 1937.
- [18] S. Sonker, A. Munjal, Sufficient conditions for triple matrices to be bounded, *Nonlinear Studies*, vol. 23, no. 4, pp. 533-542, 2016.
- [19] S. Sonker, A. Munjal, Absolute Summability Factor  $\varphi - |C, 1; \delta|_k$  of Infinite Series, *International Journal of Mathematical Analysis*, vol. 10, no. 23, pp. 1129 - 1136, 2016.
- [20] S. Sonker, A. Munjal, Approximation of the function  $f$  belong to  $Lip(\alpha, p)$  using infinite matrices of Cesàro sub-method, *Nonlinear Studies*, vol. 24, no. 1, pp. 113-125, 2017.
- [21] S. Sonker, A. Munjal, A note on boundness conditions of absolute summability  $\varphi - |A|_k$  factors, *International Conference on Advances in Science and Technology 2017*, 67, pp. 208-210. Proceedings ICASST-2017 Type A, ISBN: 9789386171429.
- [22] S. Sonker, A. Munjal, Absolute Summability  $\varphi - |C, \alpha, \beta; \delta|_k$  of Infinite Series, *Journal of Inequalities and Applications*, 2017: 168, DOI: 10.1186/s13660-017-1445-5, pp. 1-7, 2017.
- [23] S. Sonker, A. Munjal, Absolute Summability Factor  $|N, p_n|_k$  of Improper Integrals, *International Journal of Engineering and Technology*, vol. 9, no. 3S, pp. 457-462, 2017.
- [24] S. Sonker, A. Munjal, Absolute Nörlund Summability  $|N, p_n|_k$  of Improper Integrals, *National Conference on Recent Advances in Mechanical Engineering (NCRAME-2017)*, Volume-II, 90, pp. 413-415, ISBN: 978-93-86256-89-8, 2017.
- [25] S. Sonker, Xh. Z. Krasniqi, A. Munjal, A note on absolute Cesàro  $\varphi - |C, 1; \delta; l|_k$  summability factor, *International Journal of Analysis and Applications*, vol. 15, no. 1, pp. 108-113, 2017.